

Parental use of 'cry it out' in infants: no adverse effects on attachment and behavioural development at 18 months

Ayten Bilgin,¹ and Dieter Wolke^{1,2}

¹Department of Psychology, University of Warwick, Coventry, UK; ²Division of Health Sciences, Warwick Medical School, University of Warwick, Coventry, UK

Background: Leaving infant to cry it out has been the subject of discussion among researchers and parents. Nevertheless, there is paucity of empirical research investigating the association between leaving infant to cry it out and consequent crying duration and frequency, mother–infant attachment and behavioural development. **Methods:** The sample with complete longitudinal data comprised 178 infants and their caretakers. Parental use of 'leaving infant to cry out' and cry duration were assessed with maternal report at term, 3, 6 and 18 months, and frequency of crying was assessed at term, 3 and 18 months of age. Attachment was measured at 18 months using the strange situation procedure. Behavioural development of the infant was assessed with two observational measures and a parent-report questionnaire at 18 months. **Results:** The use of 'leaving infant to crying' was rare at term and increased over the next 18 months. 'Leaving infants to cry it out' at term was associated with a decrease in crying frequency at 3 months. Furthermore, leaving infants to cry it out a few times at term and often at 3 months was associated with shorter fuss/cry duration at 18 months of age. No adverse impacts of leaving infants to cry it out in the first 6 months on infant–mother attachment and behavioural development at 18 months were found. **Conclusions:** Contemporary practice of some parents in the United Kingdom to occasionally or often 'leaving infant to cry it out' during the first 6 months was not associated with adverse behavioural development and attachment at 18 months. Increased use of 'leaving to cry it out' with age may indicate differential responding by parents related to infant self-regulation. **Keywords:** Crying; cry-it-out; attachment.

Introduction

Should you leave an infant to cry it out or respond immediately and attempt to soothe him/her? This has been the subject of vehement discussion between attachment theorists and behaviourists since half a century ago (Bell & Ainsworth, 1972; Gewirtz & Boyd, 1977) and is a perennial concern for both researchers and parents (Blunden, Ether-ton, & Hauck, 2016; Ramos & Youngclarke, 2006). Attachment theorists recommended mothers to provide prompt response to infant crying according to infants' needs. On the other hand, they have advised against leaving infants to cry it out due to having concerns about its potential negative impact on the development of infant–mother attachment (Bell & Ainsworth, 1972). In other words, they have suggested that immediate responsiveness reflects sensitive parenting (Bell & Ainsworth, 1972). Indeed, there was support for this claim in an early small-sample ($N = 26$) naturalistic study conducted by Bell and Ainsworth (1972). In this study, the number of crying episodes that is ignored by the mother, the duration of maternal unresponsiveness and duration and frequency of the infant crying were observed at home during the first year at each of the four quarters for approximately 4 hr during each visit. Findings of this study showed that as the

number of ignored crying episodes increased, the frequency of crying also increased. Furthermore, it was shown that when the duration to respond to infant's crying signal increased, the duration of infant crying increased both concurrently and prospectively (Bell & Ainsworth, 1972). It was further shown that increased crying at 3 months was associated with insecure attachment at 12 months (Ainsworth, Blehar, Waters, & Wall, 1978a, 1978b). In contrast, pure behaviourists argued that leaving infants to cry it out would reduce the duration of crying, whereas immediate responding would reinforce the crying behaviour (Gewirtz & Boyd, 1977).

Surprisingly, despite it being an old controversy, there have not been many attempts to replicate the findings of Bell and Ainsworth's (1972) study. A few early attempts revealed support for the findings of Bell and Ainsworth's (1972) study; however, they had some methodological shortcomings such as failure to measure maternal unresponsiveness, short follow-up periods (i.e. 3 months) or reporting on cross-sectional associations (Belsky, Rovine, & Taylor, 1984; Crockenberg & Smith, 1982; Grossmann, Grossmann, Spangler, Suess, & Unzner, 1985). In the only complete replication study (van IJzendoorn & Hubbard, 2000), a sample of 50 mother–infant dyads were recruited to examine the association between frequency (i.e. percentage of infant crying ignored by the mother), and duration of maternal

Conflict of interest statement: No conflicts declared.

unresponsiveness (i.e. percentage of the time infant cried before maternal response) and consequent infant crying duration and frequency over a 9-month period. The study further explored the association between frequency and duration of ignoring infant crying and attachment classification at 15 months. Findings revealed that the frequency of ignoring infant's crying signals during the first 9 weeks particularly was associated with a decrease in the frequency of infant crying 9 weeks later. However, there were no significant associations between duration of infant crying before maternal response and later crying duration of the infant. Furthermore, frequency and duration of infant crying ignored by the mother did not differentiate infant–mother attachment classifications prospectively. Thus, waiting longer to respond to crying may allow the infant to make attempts of self-soothing before responding. This is contrary to the assertion by early attachment theorists that immediate responding should reduce crying (Bell & Ainsworth, 1972) and partly supportive of a behavioural control interpretation (Gewirtz & Boyd, 1977). Therefore, the existing two studies provided contradictory findings.

The current study has two objectives: First, to examine the association between the frequency of parents leaving their infant to cry it out anytime during the day at term, 3 and 6 months and cry duration and frequency during infancy at subsequent assessment points (i.e. 3, 6 or 18 months) and second, to investigate the association between frequency that parents leave their infant to cry it out during the first 6 months of infancy and attachment type and behavioural outcomes both directly observed or reported by the mother (i.e. social referencing, easiness, task persistence, aggression, poor attention/hyperactivity) at 18 months of age.

Methods

Participants

Participants were recruited at three hospitals in the East of England when they were born. The infants were assessed longitudinally at term, 3, 6 and 18 months. The sample of this study with complete longitudinal data comprised 73 very preterm/very low birthweight (VP/VLBW (<32 weeks gestation or <1,500 g birth weight) and 105 full-term infants and their mothers. Previous studies revealed no difference between the VP/VLBW and full-term samples in crying problems at 3 and 6 months and maternal sensitivity (Bilgin & Wolke, 2015, 2016). Thus, VP/VLBW and full-term samples were combined to allow for sufficient statistical power while controlling for any effect of preterm birth in all analyses. There were 101 (56.7%) male and 77 (43.3%) female participants. The mean gestational age was 35.03 ($SD = 4.91$) weeks, and the mean birth weight was 2408.98 ($SD = 1061.81$) grams. Forty-one per cent of infant participants had no siblings, 39% had one sibling, and the remaining 20% participants had more than one sibling. Furthermore, 40% had low to moderate income (yearly income of <£25 k). Mean maternal age was 30.6 ($SD = 5.82$), and 34.5% of the infants' mothers had education for more than 10 years (Table 1).

Measures

Leaving infants to cry it out. Mothers were asked to report on the frequency of 'Leaving their infant to cry it out' with the following question at term, 3, 6 and 18 months: 'Have you ever tried leaving your baby to cry it out during this time?' The answers were as follows: never, once, a few times and often. It was decided to combine the response options of 'never' and 'once' into one group, which comprised 'no cry-it-out' group since few responded leaving their infant to cry it out 'once' (22 (12.8%) at term, 9 (5.1%) at 3 months and 16 (9%) at 6 months). Thus, the analysis included three groups: no cry-it-out, a few times and often.

Crying duration and crying frequency. Mothers were asked to report how long their infant fussed/cried during the morning, afternoon, evening and at night in minutes yesterday at term, 3, 6 and 18 months and the number of bouts of fussing/crying during morning, afternoon, evening and night yesterday at term, 3 and 18 months using the Crying Pattern Questionnaire (St James-Roberts & Halil, 1991). Durations and bouts of fussing/crying during morning, afternoon, evening and night were summed to create overall cry duration and frequency of crying measure. The validity of this questionnaire has been shown to be moderate to good ($r = .51-.58$) for cry duration and moderate for fuss/cry bouts ($r = .27-.51$) when compared with a standard fuss/cry seven day diary (Wolke, Meyer, & Gray, 1994).

Attachment. Attachment type was assessed at 18 months with the strange situation procedure (SSP), a widely used and well-validated laboratory procedure to measure the quality of attachment (Ainsworth et al., 1978a, 1978b). The coders were blind to child and family characteristics and infant crying history. A third (32%) of the tapes were randomly selected for inter-rater reliability assessment, which was found to be acceptable ($\kappa = .76$). A categorical variable was created to measure attachment insecurity: 0 = secure versus 1 = insecure (insecure-avoidant and insecure-resistant).

Attachment disorganization scores were coded according to Main and Solomon's (1990) continuous scale, and a categorical variable was created: 0 = organized versus (<5) 1 = disorganized (≥ 6). Please see Appendix S1 for full description.

Observer-rated behaviour. Infant behaviour was assessed with two different observational measures. The first assessment was the Play Observation Scheme and Emotion Rating (POSER) at 18 months (Bilgin & Wolke, 2017; Wolke, 1986). POSER is an observational measure to rate maternal and infant behaviours, which includes play with a toy and free play, each lasting 2.5 min. Scales in both sessions were rated by two independent researchers who were blind to child characteristics. Each episode was viewed by the researchers a minimum of three times, focusing first on maternal behaviours, followed by infant behaviours and mother–infant joint behaviours. Overall, the coding procedure took approximately half an hour per infant–mother dyad. Infant behaviours were rated on two subscales using 9-point Likert scales (1 = very low, 9 = very high). First, activity, intensity and persistence/attentiveness were combined to a scale of poor attention/hyperactivity. Internal consistency was found to be moderate ($\alpha = .71$), and the inter-rater reliability was found to be $\kappa = .90$. Second, social referencing, emotional tone and amount of vocalization were combined to an overall scale of social referencing, that is showing positive emotional expression, vocalization and integrating mothers' suggestions in behaviour ($\alpha = .70$). The inter-rater reliability was found to be $\kappa = .91$.

The second assessment is the Tester's Rating of Infant Behaviour (TRIB) completed by a trained examiner during the Bayley Scales assessment (Jaekel, Wolke, & Bartmann, 2013;

Table 1 Demographic characteristics

Demographic characteristics	Very preterm	Full-term	Total sample
Infant sex: <i>N</i> (%)			
Female	32 (43.8%)	45 (42.9%)	77 (43.3%)
Male	41 (56.2%)	60 (57.1%)	101 (56.7%)
Gestational age (weeks): <i>M</i> (<i>SD</i>)	29.42 (1.65)	38.94 (2.25)	35.03 (4.91)
Birthweight (g): <i>M</i> (<i>SD</i>)	1285.75 (345.73)	3205.06 (554.13)	2408.98 (1061.81)
Birth order: <i>N</i> (%)			
Firstborn	31 (42.5%)	39 (37.1%)	70 (39.3%)
Maternal education			
≤10 years	45 (66.2%)	63 (64.9%)	108 (65.5%)
>10 years	23 (33.8%)	34 (35.1%)	57 (34.5%)
Maternal age: <i>M</i> (<i>SD</i>)	30.5 (5.7)	30.7 (5.9)	30.6 (5.8)
Income: <i>N</i> (%)			
£0–£25 k	30 (45.5%)	38 (36.5%)	68 (40%)
>£25 k	36 (54.5%)	66 (63.5%)	102 (60%)
Breastfeeding: <i>N</i> (%)			
Term	21 (28.8%)	55 (52.4%)	76 (42.7%)
3 Months	13 (17.8%)	31 (29.8%)	44 (24.9%)
6 Months	10 (14.3%)	13 (13.4%)	23 (13.8%)
Sleeping in parents' room: <i>N</i> (%)			
Term	54 (74.0%)	76 (72.4%)	130 (73.0%)
3 Months	43 (58.9%)	60 (57.1%)	103 (57.9%)
6 Months	23 (31.5%)	0 (0.0%)	23 (12.9%)

Wolke, Skuse, & Mathisen, 1990) at 18 months. Observations lasted on average 45 min, and 20% of the assessments were videotaped and rated by independent examiners for reliability assessment. Behaviours were rated on a 9-point Likert scale ranging from 1 = very low to 9 = very high. The six rating scales attentiveness, competence, cooperativeness, robustness/endurance, low demandingness and difficultness were combined to an overall scale of task persistence with high internal consistency ($\alpha = .95$) and $\kappa = .93$ inter-rater reliability. Ratings on adaptability, initial approach to examiner and emotional tone were combined to a total scale labelled easiness, which had high internal consistency ($\alpha = .87$) and $\kappa = .90$ inter-rater reliability.

Parent-rated behaviour problems. Parents rated the behaviour of their 18-month-old infants using the Child Behaviour and Health Questionnaire (CBHQ). The questionnaire consists of 35 items adapted from three validated scales for older children: the Strengths and Difficulties Questionnaire (SDQ; Goodman, 1997), ADHD Rating Scales (DuPaul, Power, Anastopoulos, & Reid, 1998) and the Child Behaviour Checklist (CBCL; Achenbach, 1991). The two scales derived from principal component analysis were poor attention/hyperactivity and aggression (see for details – Table S1) and had good internal consistency ($\alpha_{\text{poor attention/hyperactivity}} = .84$; $\alpha_{\text{aggression}} = .74$).

Maternal sensitivity. Maternal sensitivity was observed at 3 and 18 months of age. At 3 months of age, maternal sensitivity was measured with the Mother-Infant Structured Play Assessment (MISPA; Bilgin & Wolke, 2017; Wolke, 1999) during 2 min of play with a toy and 2 min of free play. Maternal sensitivity was coded using a 5-point scale of maternal positive emotion expression, sensitivity and stimulation adapted from three interaction coding schemes: the Play Observation Scheme and Emotion Ratings: POSER (Wolke, 1986); the Emotional Availability Scales: EAS (Biringen, Robinson, & Emde, 1993); and the Infant and Caregiver Engagement Phases: ICEP (Weinberg & Tronick, 1998). The inter-rater reliability scores for each item were high ($\kappa = .76$, $\kappa = .76$, $\kappa = .78$), and the overall internal consistency of the maternal sensitivity factor was moderate ($\alpha = .73$).

At 18 months, maternal sensitivity was observed with the POSER (Bilgin & Wolke, 2017; Wolke, 1986), the same mother-child interaction observation we used to rate infant

behaviour (see above). The maternal sensitivity factor consisted of maternal positive emotion expression, sensitivity and appropriateness of play each rated on a 9-point Likert scale (1 = low; 9 = high). The inter-rater reliability of each of the maternal behaviour items ($\kappa_{\text{positive emotion}} = .93$, $\kappa_{\text{sensitivity}} = .90$, $\kappa_{\text{appropriateness of play}} = .91$) was high. The ratings on the three items during the unstructured and structured play situation were totalled for an overall maternal sensitivity score, which had high internal consistency ($\alpha = .90$).

Statistical analysis

Analyses were performed using SPSS version 23. Logistic regression and multiple regression analyses were used to analyse the association between leaving infant to cry it out at term, 3 and 6 months and the outcome variables which were (a) crying duration and frequency of crying at the subsequent assessment point (3, 6, 18 months); (b) attachment (insecure or disorganized), observer-rated behavioural development (poor attention/hyperactivity, social referencing (POSER) and task persistence, easiness (TRIB) and parent-rated behavioural outcomes (poor attention/hyperactivity and aggression) at 18 months. Differences in sample characteristics across the three groups of 'leaving infant to cry it out' (0 = Never, 1 = A few times, 2 = Often) were analysed using chi-square tests and one-way ANOVAs. We applied Bonferroni correction to account for multiple comparisons.

Income (0 = £0–£25 k, 1 = >£25 k), preterm birth (0 = Full-Term infants; FT, 1 = Very Preterm/Very Low Birthweight infants; VP/VLBW), being firstborn and crying duration or frequency at the previous assessment point were included as covariates in all analyses. Sensitivity analysis was performed, where all analyses were repeated excluding very preterm born participants. Statistical significance was defined as $p < .05$.

Results

Characteristics of participants according to leaving infant to cry it out

Figure 1 shows the frequency of leaving infant to cry it out during the first 18 months. The majority of the

parents reported never leaving their infants to cry it out at term ($N = 109$, 63.4%), although as time went on more parents reported leaving infant to cry it out a few times ($N = 50$, 29.1% at term; $N = 87$, 48.9% at 3 months; $N = 87$, 52.1% at 6 months; $N = 65$, 36.9% at 18 months) or often ($N = 13$, 7.6% at term; $N = 23$, 12.9% at 3 months; $N = 13$, 7.8% at 6 months and $N = 55$, 31.3% at 18 months).

High-income mothers most frequently never left their infant to cry it out at term (69.8%) and 3 months (44.1%; Table 2). At 6 months, mothers who never left their infant to cry it out were older ($M = 32.6$, $SD = 4.7$) in comparison with those who left their infant to cry it out often ($M = 25.2$, $SD = 5.8$). At 18 months, mothers who left their infant to cry it out more often had higher maternal sensitivity ($M = 6.42$, $SD = 1.46$) in comparison with those who left their infant to cry it out a few times ($M = 5.74$, $SD = 1.30$), while there were no differences in maternal sensitivity during play interaction at 3 months (Table 2). Table S2 shows the bivariate correlations for all study variables.

Associations between leaving infant to cry it out and cry duration and frequency during infancy

Leaving infant to cry it out a few times at term was significantly negatively associated with crying duration at 18 months ($\beta = -.22$, $p = .004$). Furthermore, leaving infant to cry it out often at 3 months had significant negative association with crying duration at 18 months ($\beta = -.17$, $p = .03$). Otherwise, there were no significant associations between leaving infant to cry it out and crying duration at earlier assessments (Table 3).

Leaving infant to cry it out a few times ($\beta = -.17$, $p = .03$) or often ($\beta = -.18$, $p = .02$) at term was associated with frequency of crying at 3 months. No other significant associations were found between leaving infant to cry it out and the frequency of crying (Table 3). When the p -value was adjusted for

multiple comparisons using Bonferroni correction ($.05/8 = .006$), only the association between leaving infant to cry it out a few times at term and crying duration at 18 months remained significant ($\beta = -.22$, $p = .004$). When we repeated the analyses excluding very preterm infants, findings remained the same (Table S3a).

Associations between leaving infant to cry it out at term, 3 and 6 months and attachment at 18 months

There were no significant associations between leaving infant to cry it out during the first 6 months and insecure or disorganized attachment at 18 months (Table 4). When we excluded very preterm infants, findings remained the same (Table S3b).

Associations between leaving infant to cry out at term, 3 and 6 months and behavioural development at 18 months

Table 4 shows that there were no significant associations between leaving infant to cry it out during the first 6 months and behavioural development at 18 months as observed during a play situation (POSER) or a cognitive assessment (TRIB). Similarly, there were no significant associations between leaving infant to cry it out during the first 6 months and parent-rated behaviour at 18 months (Table 4). When we excluded very preterm infants and the p -value was adjusted for multiple comparisons, findings remained the same (Table S3b).

Discussion

This study found, first, that frequency of leaving infants to cry it out in the first 6 months in infancy is not associated with an increase in crying duration or frequency of crying up to 18 months of age. Rather, cry duration was found to be lower at 18 months if




Figure 1 Frequency of leaving infant to cry out at term, 3, 6 and 18 months

Table 2 Characteristics of mothers and infants according to response to crying at term, 3, 6 and 18 months

	Total number	Leaving infant to cry it out			<i>p</i>
		Never	A few times	Often	
Term					
Maternal education					
≤10 years	104	63 (60.6%)	30 (28.8%)	11 (10.6%)	.22
>10 years	56	40 (71.4%)	14 (25%)	2 (3.6%)	
Maternal age	172	31.3 (5.3)	29.7 (6.6)	27.9 (6.4)	.07
Income					
£0–£25 k	68	37 (54.4%)	21 (30.9%)	10 (14.7%)	.01
>£25 k	96	67 (69.8%)	27 (28.1%)	2 (2.1%)	
Infant sex					
Female	75	67 (69.1%)	23 (23.7%)	7 (7.2%)	.19
Male	97	42 (56%)	27 (36%)	6 (8%)	
Preterm birth					
Very preterm	67	38 (56.7%)	21 (31.3%)	8 (11.9%)	.15
Full-term	105	71 (67.6%)	29 (27.6%)	5 (4.8%)	
Firstborn	67	46 (68.7%)	18 (26.9%)	3 (4.5%)	.36
Breastfeeding					
Yes	74	51 (68.9%)	20 (27%)	3 (4.1%)	.23
No	98	58 (59.2%)	30 (30.6%)	10 (10.2%)	
Sleeping at parents' room					
Yes	130	82 (63.1%)	37 (28.5%)	11 (8.5%)	.72
No	42	27 (64.3%)	13 (31%)	2 (4.8%)	
3 Months					
Maternal education					
≤10 years	108	38 (35.2%)	52 (48.1%)	18 (16.7%)	.10
>10 years	57	25 (43.9%)	29 (50.9%)	3 (5.3%)	
Maternal age	178	31.5 (5.5)	30.3 (5.9)	28.9 (6.2)	.14
Income					
£0–£25 k	68	20 (29.4%)	34 (50%)	14 (20.6%)	.04
>£40 k	102	45 (44.1%)	48 (47.1%)	9 (8.8%)	
Maternal sensitivity	174	3.96 (0.51)	3.90 (0.61)	3.80 (0.51)	.49
Infant sex					
Female	77	29 (37.7%)	36 (46.8%)	12 (15.6%)	.64
Male	101	39 (38.6%)	51 (50.5%)	11 (10.9%)	
Preterm birth					
Very preterm	73	26 (35.6%)	32 (43.8%)	15 (20.5%)	.04
Full-term	105	42 (40%)	55 (52.4%)	8 (7.6%)	
Firstborn	70	26 (37.1%)	34 (48.6%)	10 (14.3%)	.91
Breastfeeding					
Yes	44	18 (40.9%)	22 (50%)	4 (9.1%)	.66
No	133	49 (36.8%)	65 (48.9%)	19 (14.3%)	
Sleeping at parents' room					
Yes	103	38 (36.9%)	55 (53.4%)	10 (9.7%)	.21
No	75	30 (40%)	32 (42.7%)	13 (17.3%)	
6 Months					
Maternal education					
≤10 years	102	37 (36.3%)	54 (52.9%)	11 (10.8%)	.16
>10 years	53	26 (49.1%)	25 (47.2%)	2 (3.8%)	
Maternal age	167	32.6 (4.7)	30.3 (5.7)	25.2 (5.8)	<.001
Income					
£0–£40k	63	21 (33.3%)	33 (52.4%)	9 (14.3%)	.06
>£40 k	97	42 (43.3%)	51 (52.6%)	4 (4.1%)	
Infant sex					
Female	71	23 (32.4%)	40 (56.3%)	8 (11.3%)	.12
Male	96	44 (45.8%)	47 (49%)	5 (5.2%)	
Preterm birth					
Very preterm	70	30 (42.9%)	33 (47.1%)	7 (10%)	.46
Full-term	97	37 (38.1%)	54 (55.7%)	6 (6.2%)	
Firstborn	61	20 (32.8%)	37 (60.7%)	4 (6.6%)	.24
Breastfeeding					
Yes	23	10 (43.5%)	12 (52.2%)	1 (4.3%)	.79
No	144	57 (39.6%)	75 (52.1%)	12 (8.3%)	
Sleeping at parents' room					
Yes	23	13 (56.5%)	9 (39.1%)	1 (4.3%)	.22
No	144	54 (37.5%)	78 (54.2%)	12 (8.3%)	

(continued)

Table 2 (continued)

	Total number	Leaving infant to cry it out			<i>p</i>
		Never	A few times	Often	
18 Months					
Maternal education					
≤10 years	107	35 (32.7%)	38 (35.5%)	34 (31.8%)	.51
>10 years	56	15 (26.8%)	25 (44.6%)	16 (28.6%)	
Maternal age	176	30.4 (4.9)	31.2 (6.2)	30.4 (6.1)	.64
Income					
£0–£40 k	68	24 (35.3%)	25 (36.8%)	19 (27.9%)	.57
>£40 k	102	29 (28.4%)	38 (37.3%)	35 (34.3%)	
Maternal sensitivity	166	5.82 (1.46)	5.74 (1.30)	6.42 (1.46)	.03
Infant sex					
Female	77	24 (31.2%)	29 (37.7%)	24 (31.2%)	.98
Male	99	32 (32.3%)	36 (36.4%)	31 (31.3%)	
Preterm birth					
Very preterm	71	27 (38%)	31 (43.7%)	13 (18.3%)	.01
Full-term	105	29 (27.2%)	34 (32.4%)	42 (40%)	
Firstborn	70	22 (31.4%)	28 (40%)	20 (28.6%)	.75

Please note that boldface type indicates significant differences.

Table 3 Predicting cry duration and cry frequency during infancy according to frequency of leaving infant to cry out

	Cry duration at 3 months		Cry duration at 6 months		Cry duration at 18 months		Cry frequency at 3 months		Cry frequency at 18 months	
	β	<i>p</i>	β	<i>p</i>	β	<i>p</i>	β	<i>p</i>	β	<i>p</i>
Leaving infant to cry it out at term										
Never	[Reference]		[Reference]		[Reference]		[Reference]		[Reference]	
A few times	-.08	.24	-.01	.89	-.22	.004	-.17	.03	-.03	.69
Often	.07	.35	-.05	.56	.05	.54	-.18	.02	-.04	.61
Leaving infant to cry it out at 3 months										
Never			[Reference]		[Reference]				[Reference]	
A few times			.01	.88	-.03	.74			.13	.09
Often			-.05	.51	-.17	.03			-.10	.24
Leaving infant to cry it out at 6 months										
Never					[Reference]				[Reference]	
A few times					.15	.06			.14	.09
Often					-.07	.36			-.11	.18

Please note that the analysis was controlled for income, preterm birth, firstborn and cry duration in the previous assessment for each regression. Boldface type indicates significant associations.

parents left their infants to cry it out for a few times at term and often at 3 months. Furthermore, leaving infants to cry it out at term both a few times or often was negatively associated with frequency of crying at 3 months. The strength of the associations was weak. Second, frequency of leaving infants to cry it out in the first 6 months in infancy was not found to be associated with either adverse behavioural effects on infant development or infant–mother attachment at 18 months of age. Thus, whether contemporary parents respond immediately or leave their infant to cry it out a few times or often might not be associated with short- or long-term adverse effects on infant behaviour or quality of infant–mother relationship during the first 18 months in this UK sample.

This study does not provide support for predictions made by either attachment theory or learning theory. Thus, we neither recommend leaving infant to cry out nor responding immediately. Rather, our

findings are consistent with an approach to parenting that is intuitive and adapts to infant demands according to infant regulatory competencies across infancy (Papousek & Papousek, 1996). In this UK sample, most mothers appear to intuitively follow a differential responsiveness approach with most of them responding immediately at term and then starting to adopt a differential response approach from 3 months onward. Crying in the first 3 months of age is biologically driven, and patterns of crying (the crying curve) have been found to be similar across cultures and countries (Barr, 1990; Wolke, Bilgin, & Samara, 2017). Up to 40% of crying in the first 3 months of life has been consistently reported to be inconsolable independent of highly responsive ‘proximal’ care or not (Lucassen et al., 2001; St James-Roberts et al., 2006; Wolke et al., 2017). Furthermore, intervention studies found that increased maternal responsiveness to crying and

Table 4 Predicting attachment and behavioural outcomes at 18 months according to leaving infant to cry it out at term, 3 and 6 months

	Observer ratings										Parent ratings								
	Disorganized attachment					POSER					TRIB			Poor attention/hyperactivity			Aggression		
	OR	95% CI	OR	95% CI	β	p	β	p	β	p	β	p	β	p	β	p	β	p	
Leaving infants to cry it out at term																			
Never	[Reference]		[Reference]		[Reference]		[Reference]		[Reference]		[Reference]		[Reference]		[Reference]		[Reference]		[Reference]
A few times	0.83	0.39–1.78	0.61	0.26–1.47	.05	.53	-.09	.25	-.02	.78	-.02	.75	-.03	.67	-.03	.79	.02	.03	.79
Often	0.87	0.23–3.3	0.81	0.19–3.51	-.08	0.32	-.09	0.27	-.06	0.46	-.06	0.42	.08	0.36	.03	0.72			
Leaving infants to cry it out at 3 months																			
Never	[Reference]		[Reference]		[Reference]		[Reference]		[Reference]		[Reference]		[Reference]		[Reference]		[Reference]		[Reference]
A few times	1.01	0.53–1.95	0.87	0.42–1.83	.05	.50	-.12	.12	-.06	.44	-.05	.52	-.03	.72	-.03	.29	-.08	.04	.29
Often	1.19	0.46–3.10	0.66	0.21–2.07	.02	.77	-.01	.92	-.01	.94	-.01	.97	-.03	.72	-.03	.60			.60
Leaving infants to cry it out at 6 months																			
Never	[Reference]		[Reference]		[Reference]		[Reference]		[Reference]		[Reference]		[Reference]		[Reference]		[Reference]		[Reference]
A few times	1.20	0.58–2.46	0.86	0.39–1.91	.05	.52	.01	.92	-.09	.29	-.08	.32	-.05	.59	-.05	.60			.60
Often	1.12	0.31–4.10	1.10	0.27–4.80	.03	.74	-.07	.39	.05	.51	.05	.54	-.07	.43	-.07	.39			.39

Please note that the analysis was controlled for income, preterm birth, firstborn and cry duration in the previous assessment for each regression. POSER: Play Observation Scheme and Emotion Rating; TRIB: Tester's Rating of Infant Behaviour.

close physical contact (i.e. carrying of baby), in particular for colic crying, did not reduce crying behaviour (Barr et al., 1991; Douglas & Hill, 2011; St James-Roberts, Hurry, Bowyer, & Barr, 1995). Thus, converging evidence suggests that neither immediate responding nor not responding during the first 3 months – as suggested by behaviourists – alters infants' crying duration subsequently in infancy. However, there may be a small association between leaving infant to cry it out at term and frequency of crying subsequently at 3 months. Likewise, the findings of the van IJzendoorn and Hubbard (2000) study, leaving infants to cry it out a few times is associated with a decrease in frequency of crying subsequently.

Leaving an infant to cry it out might not reflect parental neglect but it may rather reflect authoritative parenting which includes both limit setting and high levels of emotional warmth (Baumrind, 1967). Delayed responding has been suggested for dealing with infant night waking signalled by infant crying from 6 months onwards. Hiscock et al. (2007) showed that intervention involving leaving infant to cry it out gradually at 7 months decreased night crying and thus sleeping problems at 10 months and 12 months. This method of leaving infant to cry it out using a graded approach corresponds with the concept of authoritative parenting (Baumrind, 1967) and is consistent with a differential responsiveness framework (van IJzendoorn & Hubbard, 2000; Wolke, 2019). Helping infants to learn how to regulate their crying behaviour is a type of limit setting and considered beneficial when combined with parental warmth (Baumrind, 1967). This is consistent with the finding here that maternal sensitivity in play situations was not or positively associated (at 18 months) with leaving infant cry it out. On the contrary, trying to respond to a crying infant immediately every time might put mothers under a lot of stress, increase fatigue (Barr et al., 2014; Long et al., 2018) and may increase depression (Petzoldt, 2018). Moreover, mothers' own expectations of responding to their crying infants immediately might even increase later infant night waking (Galbally, Watson, Teti, & Lewis, 2018). For infants' healthy development, the importance of maternal self-care and good maternal mental health is invaluable (Kurth, Kennedy, Spichiger, Hosli, & Stutz, 2011). Knowing that leaving infant to cry it out gradually decreases maternal depression (Hiscock et al., 2007), it is more likely to be beneficial for infant emotional and behavioural development in comparison with having ill effects. However, the number of studies on leaving infant to cry it out is still scarce and longitudinal studies of different cohorts are needed to confirm this finding.

The second argument put forward against leaving infants to cry it out is that it may increase the risk of insecure mother–infant attachment as well as behavioural problems (Bell & Ainsworth, 1972; Murray &

Ramchandani, 2007). The current study found no associations between leaving infants to cry it out during the first 6 months and insecure or disorganized attachment at 18 months consistent with the findings of van Ijzendoorn and Hubbard's (2000) study. Furthermore, no associations were found between frequency of leaving infants to cry it out and infant behaviour at 18 months. Similarly, sleep intervention studies and studies using timeout (Knight, Albright, Deling, Dore-Stites, & Drayton, 2019) have not shown any adverse effect of using a graded extinction approach in response to waking and crying at night on infant-mother attachment or behaviour after one year (Gradisar et al., 2016) and 5 years (Price, Wake, Ukoumunne, & Hiscock, 2012). Indeed, a majority of parents reported to be successful in adopting a behavioural approach in the community to deal with night waking and crying (Honaker, Schwichtenberg, Kreps, & Mindell, 2018). The children fell asleep faster and woke less within a week.

A further argument against leaving infants to cry it out has been the suggestion that it may increase infants' stress levels, which may be harmful to the infant (Etherton, Blunden, & Hauck, 2016; Murray & Ramchandani, 2007). However, there is no empirical evidence that leaving infants to cry it out actually induces stress to the infant. The only study on the stress levels of infants after a 5-day intervention programme including leaving infants to cry it out gradually revealed no change in infant cortisol levels (Middlemiss, Granger, Goldberg, & Nathans, 2012). Short- and long-term chronic stress impacts humans differently (Monaghan & Haussmann, 2015), and early stress may increase adaptive strategies (Ellis & Del Giudice, 2019). It has been shown that short-term stress is beneficial in terms of its positive impact on resiliency to stress in later life (Ellis & Del Giudice, 2019; Monaghan & Haussmann, 2015). Differential responding as part of intuitive parenting provides infants the opportunity to soothe themselves by waiting before intervening when crying and allows them to explore ways of self-soothing, acquiring a first sense of self and regulating their behavioural state (Wolke, 2019). If leaving infants to cry it out is occurring within the context of a warm mother-infant relationship, no adverse impacts have been demonstrated.

This prospective longitudinal study has several strengths, including a larger sample than previous studies, adjustment for several important covariates, using repeated assessment of frequency of leaving infant to cry it out during infancy, observer assessed infant-mother attachment and the assessment of behavioural outcomes using both researcher observations and parental reports. There are also limitations. First, frequency of leaving infant to cry it out and cry duration and frequency were assessed via maternal report. This assessment represents maternal perception of her own

behaviour and her infant's crying behaviour. Direct independent observation over long periods would be desirable but is less feasible in a prospective study with a good sample size and repeated measurements. Where direct observations of crying amounts and maternal report have been compared, the correlations exceeded $r = .80$ (St James-Roberts, Hurry, & Bowyer, 1993). Furthermore, maternal report has the benefit of capturing behaviours over longer time periods. Second, the correlational design of the current study does not allow to determine whether there is a causal relationship between leaving infant to cry it out and the assessed outcomes. However, randomization in controlled trials may not be possible due to strongly held views by parents. Third, frequency of crying was not assessed at 6 months. Thus, the current study did not allow us to investigate the associations between leaving infant to cry out at 3 months and frequency of crying at 6 months. Fourth, the validity information of CBHQ measure was not available. Nevertheless, this measure was formed by items from valid measures. Fifth, the current study did not allow to distinguish between leaving infant to cry out during daytime and night time. Thus, we could not distinguish whether parents may wait less or longer when intervening at day than night time. Future studies may address responsiveness to crying during daytime and night time separately. Lastly, the current study included a large group of infants who were born very preterm or very low birthweight (41%). All analyses were controlled for the role of very preterm birth. Furthermore, sensitivity analysis excluding very preterm born participants did not alter the main findings, revealing wider confidence intervals.

In conclusion, contemporary practice by some parents to occasionally or often 'leaving infant to cry it out' during the first 6 months was not associated with adverse behavioural development and attachment at 18 months. Nor was 'leaving infant to cry it out' associated with reduced maternal sensitivity in interaction at 3 or 18 months of age. Increased use of 'leaving to cry it out' with age may indicate differential responding by mothers to aid the development of infant self-regulation.

Supporting information

Additional supporting information may be found online in the Supporting Information section at the end of the article:

Appendix S1. Full description of attachment coding.

Table S1. Variables forming poor attention/hyperactivity and aggression subscales in Child Behaviour and Health Questionnaire (CBHQ).

Table S2. Bivariate correlations among study variables.

Table S3a. Predicting cry duration and cry frequency during infancy according to frequency of leaving infant to cry out.

Table S3b. Predicting attachment and behavioural outcomes at 18 months according to leaving infant to cry it out at term, 3 and 6 months.

Acknowledgements

This study was supported by grant 1590/611 from the Health Foundation, UK. The authors would like to thank the researchers who assisted in recruitment and data collection: Tina Gutbrod, Libi Rust and Karine Edme. The authors would also like to thank the participating hospitals (Addenbrookes Hospital, Cambridge; Luton and Dunstable Hospital, Luton; and

Queen Elizabeth II Hospital, Welwyn Garden City) and the parents and their children. Furthermore, the authors would like to thank Eva Liu and Dr. Lara Ayala-Nunes for providing feedback on the initial manuscript. The authors have declared that they have no competing or potential conflicts of interest.

Correspondence

Dieter Wolke, Department of Psychology, University of Warwick, Coventry CV4 7AL, UK; Email: D.Wolke@warwick.ac.uk

Key points

- It has been the subject of vehement discussion between attachment theorists and behaviourists since half a century ago whether parents should leave their infant to cry it out or respond immediately and attempt to soothe him/her.
- No adverse impacts of leaving infants to cry it out in the first 6 months on infant–mother attachment and behavioural development at 18 months were found.
- Parents may be reassured that there is no harmful impact of leaving their infants to cry it out sometimes or often during infancy while a parent is present, the infant is safe, and they monitor the infant's crying.

References

- Achenbach, T.M. (1991). *Manual for the child behavior checklist/4-18 and 1991 profile*. Burlington, VT: University of Vermont, Department of Psychiatry.
- Ainsworth, M., Blehar, M., Waters, E., & Wall, S. (1978a). *Patterns of attachment*. Hillsdale, NJ: Erlbaum.
- Ainsworth, M.D.S., Blehar, M.C., Waters, E., & Wall, S. (1978b). *Patterns of attachment: A psychological study of the strange situation*. Oxford, UK: Lawrence Erlbaum.
- Barr, R.G. (1990). The normal crying curve: What do we really know? *Developmental Medicine & Child Neurology*, 32, 356–362.
- Barr, R.G., Fairbrother, N., Pauwels, J., Green, J., Chen, M., & Brant, R. (2014). Maternal frustration, emotional and behavioural responses to prolonged infant crying. *Infant Behavior and Development*, 37, 652–664.
- Barr, R.G., McMullan, S.J., Spiess, H., Leduc, D.G., Yaremko, J., Barfield, R., ... & Hunziker, U.A. (1991). Carrying as colic “therapy”: A randomized controlled trial. *Pediatrics*, 87, 623–630.
- Baumrind, D. (1967). Child care practices anteceding three patterns of preschool behavior. *Genetic Psychology Monographs*, 75, 43–88.
- Bell, S.M., & Ainsworth, M.D. (1972). Infant crying and maternal responsiveness. *Child Development*, 43, 1171–1190.
- Belsky, J., Rovine, M., & Taylor, D.G. (1984). The Pennsylvania Infant and Family Development Project, III: The origins of individual differences in infant–mother attachment: maternal and infant contributions. *Child Development*, 55, 718–728.
- Bilgin, A., & Wolke, D. (2015). Maternal sensitivity in parenting preterm children: A meta-analysis. *Pediatrics*, 136, e177–e193.
- Bilgin, A., & Wolke, D. (2016). Regulatory problems in very preterm and full-term infants over the first 18 months. *Journal of Developmental and Behavioral Pediatrics*, 37, 298–305.
- Bilgin, A., & Wolke, D. (2017). Associations between feeding problems and maternal sensitivity across infancy: Differences in very preterm and full-term infants. *Journal of Developmental and Behavioral Pediatrics*, 38, 538–544.
- Biringen, Z., Robinson, J., & Emde, R.N. (1993). *The emotional availability scales* (2nd ed.). Fort Collins, CO: Department of Human Development and Family Studies, Colorado State University.
- Blunden, S., Etherton, H., & Hauck, Y. (2016). Resistance to cry intensive sleep intervention in young children: Are we ignoring children's cries or parental concerns? *Children*, 3, 8.
- Crockenberg, S.B., & Smith, P. (1982). Antecedents of mother–infant interaction and infant irritability in the first three months of life. *Infant Behavior and Development*, 5, 105–119.
- Douglas, P., & Hill, P. (2011). Managing infants who cry excessively in the first few months of life. *BMJ*, 343, d7772.
- DuPaul, G.J., Power, T.J., Anastopoulos, A.D., & Reid, R. (1998). *ADHD Rating Scale—IV: Checklists, norms, and clinical interpretation*. New York: Guilford Press.
- Ellis, G.B., & Del Giudice, M. (2019). Developmental adaptation to stress: An evolutionary perspective. *Annual Review of Psychology*, 70, 111–139.
- Etherton, H., Blunden, S., & Hauck, Y. (2016). Discussion of extinction-based behavioral sleep interventions for young children and reasons why parents may find them difficult. *Journal of Clinical Sleep Medicine*, 12, 1535–1543.
- Galbally, M., Watson, S.J., Teti, D., & Lewis, A.J. (2018). Perinatal maternal depression, antidepressant use and infant sleep outcomes: Exploring cross-lagged associations in a pregnancy cohort study. *Journal of Affective Disorders*, 238, 218–225.
- Gewirtz, J.L., & Boyd, E.F. (1977). Does maternal responding imply reduced infant crying? A critique of the 1972 Bell and Ainsworth report. *Child Development*, 48, 1200–1207.
- Goodman, R. (1997). The Strengths and Difficulties Questionnaire: A research note. *Journal of Child Psychology and Psychiatry*, 38, 581–586.

- Gradisar, M., Jackson, K., Spurrier, N.J., Gibson, J., Whitham, J., Williams, A.S., ... & Kennaway, D.J. (2016). Behavioral interventions for infant sleep problems: A Randomized Controlled Trial. *Pediatrics*, *137*, e20151486.
- Grossmann, K., Grossmann, K.E., Spangler, G., Suess, G., & Unzner, L. (1985). Maternal sensitivity and newborns' orientation responses as related to quality of attachment in Northern Germany. *Monographs of the Society for Research in Child Development*, *50*, 233–256.
- Hiscock, H., Bayer, J., Gold, L., Hampton, A., Ukoumunne, O.C., & Wake, M. (2007). Improving infant sleep and maternal mental health: A cluster randomised trial. *Archives of Disease in Childhood*, *92*, 952–958.
- Honaker, S.M., Schwichtenberg, A.J., Kreps, T.A., & Mindell, J.A. (2018). Real-world implementation of infant behavioral sleep interventions: Results of a parental survey. *The Journal of Pediatrics*, *199*, 106–111.e2.
- Jaekel, J., Wolke, D., & Bartmann, P. (2013). Poor attention rather than hyperactivity/impulsivity predicts academic achievement in very preterm and full-term adolescents. *Psychological Medicine*, *43*, 183–196.
- Knight, R.M., Albright, J., Deling, L., Dore-Stites, D., & Drayton, A.K. (2019). Longitudinal relationship between time-out and child emotional and behavioral functioning. *Journal of Developmental and Behavioral Pediatrics*, *41*(1), 31–37.
- Kurth, E., Kennedy, H.P., Spichiger, E., Hosli, I., & Stutz, E.Z. (2011). Crying babies, tired mothers: What do we know? A systematic review. *Midwifery*, *27*, 187–194.
- Long, J., Powell, C., Bamber, D., Garratt, R., Brown, J., Dyson, S., & James-Roberts, I.S. (2018). Development of materials to support parents whose babies cry excessively: findings and health service implications. *Primary Health Care Research & Development*, *19*, 320–332.
- Lucassen, P.L.B.J., Assendelft, W.J.J., van Eijk, J.T.M., Gubbels, J.W., Douwes, A.C., & van Geldrop, W.J. (2001). Systematic review of the occurrence of infantile colic in the community. *Archives of Disease in Childhood*, *84*, 398–403.
- Main, M., & Solomon, J. (1990). Procedures for identifying infants as disorganized/disoriented during the Ainsworth Strange Situation. In M. T. Greenberg, D. Cicchetti & E. M. Cummings (Eds.), *Attachment in the preschool years: Theory, research, and intervention* (pp. 121–160). Chicago, IL: University of Chicago Press.
- Middlemiss, W., Granger, D.A., Goldberg, W.A., & Nathans, L. (2012). Asynchrony of mother-infant hypothalamic-pituitary-adrenal axis activity following extinction of infant crying responses induced during the transition to sleep. *Early Human Development*, *88*, 227–232.
- Monaghan, P., & Haussmann, M.F. (2015). The positive and negative consequences of stressors during early life. *Early Human Development*, *91*, 643–647.
- Murray, L., & Ramchandani, P. (2007). Might prevention be better than cure? *Archives of Disease in Childhood*, *92*, 943–945.
- Papousek, M., & Papousek, H. (1996). Infant colic, state regulation, and interaction with parents: A systems approach. In M.H.B.J.L. Genevro (Ed.), *Child development and behavioral pediatrics: Toward understanding children and health* (pp. 11–33). Hillsdale, NJ: Lawrence Erlbaum.
- Petzoldt, J. (2018). Systematic review on maternal depression versus anxiety in relation to excessive infant crying: It is all about the timing. *Archives of Women's Mental Health*, *21*, 15–30.
- Price, A.M.H., Wake, M., Ukoumunne, O.C., & Hiscock, H. (2012). Five-year follow-up of harms and benefits of behavioral infant sleep intervention: Randomized Trial. *Pediatrics*, *130*, 643–651.
- Ramos, K.D., & Youngclarke, D.M. (2006). Parenting advice books about child sleep: cosleeping and crying it out. *Sleep*, *29*, 1616–1623.
- St James-Roberts, I., Alvarez, M., Csipke, E., Abramsky, T., Goodwin, J., & Sorgenfrei, E. (2006). Infant crying and sleeping in London, Copenhagen and when parents adopt a “proximal” form of care. *Pediatrics*, *117*(6), e1146–e1155.
- St James-Roberts, I., & Halil, T. (1991). Infant crying patterns in the first year: Normal community and clinical findings. *Journal of Child Psychology and Psychiatry*, *32*, 951–968.
- St James-Roberts, I., Hurry, J., & Bowyer, J. (1993). Objective confirmation of crying durations in infants referred for excessive crying. *Archives of Disease in Childhood*, *68*, 82–84.
- St James-Roberts, I., Hurry, J., Bowyer, J., & Barr, R.G. (1995). Supplementary carrying compared with advice to increase responsive parenting as interventions to prevent persistent infant crying. *Pediatrics*, *95*, 381–388.
- van IJzendoorn, M.H., & Hubbard, F.O. (2000). Are infant crying and maternal responsiveness during the first year related to infant-mother attachment at 15 months? *Attachment & Human Development*, *2*, 371–391.
- Weinberg, M.K., & Tronick, E.Z. (1998). *Infant and caregiver engagement phases system*. Boston: Children's Hospital and Harvard Medical School.
- Wolke, D. (1986). *Play observation scheme and emotion rating*. Hatfield, UK: University of Hertfordshire.
- Wolke, D. (1999). *The mother-infant structured play assessment (MISPA)*. Hatfield, UK: University of Hertfordshire.
- Wolke, D. (2019). Persistence of infant crying, sleeping and feeding problems: Need for prevention. *Archives of Disease in Childhood*, *104*, 1022–1023.
- Wolke, D., Bilgin, A., & Samara, M. (2017). Systematic review and meta-analysis: Fussing and crying durations and prevalence of colic in infants. *Journal of Pediatrics*, *185*, 55–61.e54.
- Wolke, D., Meyer, R., & Gray, P. (1994). Validity of the Crying Pattern Questionnaire in a sample of excessively crying babies. *Journal of Reproductive and Infant Psychology*, *12*, 105–114.
- Wolke, D., Skuse, D., & Mathisen, B. (1990). Behavioral style in failure-to-thrive infants - a preliminary communication. *Journal of Pediatric Psychology*, *15*, 237–254.

Accepted for publication: 22 January 2020